

## HA-Tag Rabbit pAb

www.dia-an.cn

Cat#: 3063

027-59325199


sevice@dia-an.com

Only For Research. Not For Diagnosis.


<b>Synonyms:</b>	YPYDVPDYA
<b>Attribute:</b>	Rabbit Polyclonal Antibody
<b>Isotype:</b>	IgG
<b>Purity:</b>	Antigen Affinity Purification
<b>Application:</b>	ELISA, WB, IP, IF
<b>Calculated MW:</b>	
<b>Observed MW:</b>	
<b>Reactivity:</b>	Human,Mouse,Rat,Zebra fish
<b>Immunogen:</b>	Peptide YPYDVPDYA
<b>Buffer:</b>	PBS with 0.1% sodium azide and 50% glycerol, pH 7.2
<b>Storage:</b>	Store at -20°C. Do not aliquot
<b>Recommended Dilution:</b>	WB: 1:2000-5000 IP: 3ug/sample IF: 1:800-1600

### Background:

The HA tag is derived from the HA-molecule corresponding to amino acids 98-106. It has been extensively used as a general epitope tag in expression vectors. Many recombinant proteins have been engineered to express the HA tag, which does not appear to interfere with the bioactivity or the biodistribution of the recombinant protein.



Western Blotting of HA-Tag fused recombinant protein with Anti-HA Rabbit Polyclonal antibody at dilution of 1:1000



Immunofluorescent analysis of HA-GFP transfected 293 cells with Anti-HA Rabbit 3063 at dilution of 1:800

We focus on precise protein quantification