

HA-Tag Rabbit mAb

www.dia-an.cn

Cat#: 3063R

027-59325199


sevice@dia-an.com

Only For Research. Not For Diagnosis.


Synonyms:	YPYDVPDYA
Attribute:	Rabbit Monoclonal Antibody
Isotype:	IgG
Purity:	Antigen Affinity Purification
Application:	ELISA, WB, IP, IF
Reactivity:	Protein fused with HA tag
Immunogen:	Peptide YPYDVPDYA
Buffer:	PBS with 0.1% sodium azide and 50% glycerol, pH 7.2
Storage:	Store at -20°C. Do not aliquot
Recommended Dilution:	WB: 1:2000-1:10,000 IP: 3ug/sample IF: 1:100-1:200

Background:

The HA tag is derived from the HA-molecule corresponding to amino acids 98-106. It has been extensively used as a general epitope tag in expression vectors. Many recombinant proteins have been engineered to express the HA tag, which does not appear to interfere with the bioactivity or the biodistribution of the recombinant protein.



Western Blotting of HA-Tag fused recombinant protein with Anti-HA Rabbit Monoclonal Antibody at dilution of 1:5000



Immunofluorescent analysis of 293F cells transfected with the HA-GFP, using 3063R at 1:2000 dilution

Western blot of Zebra fish whole lysates with anti-H2A Rabbit Polyclonal Antibody at dilution of 1:1000

We focus on precise protein quantification

